



Nitrogen as a factor for enhanced carbon sequestration

Results from four NitroEurope-IP forest supersites.

NitroEurope-IP Forest Synthesis Team

Published in:
Geophysical Research Abstracts

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
NitroEurope-IP Forest Synthesis Team (2012). Nitrogen as a factor for enhanced carbon sequestration: Results from four NitroEurope-IP forest supersites. *Geophysical Research Abstracts*, 14, EGU2012-7950.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Nitrogen as a factor for enhanced carbon sequestration: Results from four NitroEurope-IP forest supersites

A. Ibrom and the NitroEurope-IP Forest Synthesis Team

Department of Chemical and Biochemical Engineering, DTU, Lyngby / Roskilde, Denmark

Nitrogen (N) fertilization, both intended and unintended, interacts with carbon cycling in terrestrial ecosystems, because the major processes of carbon (C) turnover depend on enzymes and thus on N availability. Comparisons between annual carbon dioxide flux (CO_2) budgets and wet N deposition in forests showed a very strong linear increase of CO_2 sequestration with increased N deposition. After considering total rather than only wet N deposition the ratios between increased carbon uptake and atmospheric N input were closer to C/N that can be found in wood. This suggested that the observed ecosystems responses to enhanced N inputs were mainly driven by plant responses. Finally, looking at changes in soil organic matter changes indicated even lower sensitivities of carbon sequestration to N addition. The objective of this study is to describe the mechanisms of the responses and the fate of the N in the ecosystem based on results from intensively investigated forest sites.

Within the European NitroEurope-IP project the annual fluxes and pool sizes of C and N were estimated in four so-called forest supersites, including temperate coniferous forests in Southern Germany (Höglwald) and in the Netherlands (Speulderbos), one temperate beech forest close to Sorø on Zealand in Denmark and a boreal pine forest (Hyytiälä, Southern Finland). Due to differences in vegetation, bedrock and climate history, soils differed in acidity, organic matter content and biological activity; the levels of atmospheric N deposition varied from very low (Hyytiälä) to high (the other sites).

Comparisons of N and C budgets of plants and soils confirmed a simple and stoichiometric effect $dC_{\text{uptake}}/dN_{\text{dep}} = \text{constant}$ and in the order of magnitude of $(\text{C/N})_{\text{wood}}$ for plants but not for soils and thus not for the forest ecosystems as a whole. Differences in soil processes as indicated by the differing C/N of SOM, differing amounts of N stored in the soil and considerable differences in N leaching rates even at comparable N deposition levels, showed clearly that the diversity of soils play a large role in the N use for C sequestration and thus for the beneficial effects of additional N loads on climate change mitigation effects in forests. An important conclusion of the study for intended forest fertilization is to consider N leaching to the ground water, which might even enhance the greenhouse effect through increased N_2O emissions from streams, estuaries and coasts rather than mitigating it via increased CO_2 sequestration at the forest site.

Acknowledgements

This work has been funded by the European Commission via the NitroEurope and CarboEurope integrated projects.